

Guidelines for Management of Incidental Pulmonary Nodules Detected on CT Images: From the Fleischner Society 2017.

MacMahon H1, Naidich DP1, Goo JM1, Lee KS1, Leung ANC1, Mayo JR1, Mehta AC1, Ohno Y1, Powell CA1, Prokop M1, Rubin GD1, Schaefer-Prokop CM1, Travis WD1, Van Schil PE1, Bankier AA1.

Radiology. 2017 Jul;284(1):228-243. doi: 10.1148/radiol.2017161659. Epub 2017 Feb 23.

<https://www.ncbi.nlm.nih.gov/pubmed/28240562>

A: Solid Nodules*				
Nodule Type	Size			Comments
	<6 mm (<100 mm ³)	6–8 mm (100–250 mm ³)	>8 mm (>250 mm ³)	
Single				
Low risk [†]	No routine follow-up	CT at 6–12 months, then consider CT at 18–24 months	Consider CT at 3 months, PET/CT, or tissue sampling	Nodules <6 mm do not require routine follow-up in low-risk patients (recommendation 1A).
High risk [†]	Optional CT at 12 months	CT at 6–12 months, then CT at 18–24 months	Consider CT at 3 months, PET/CT, or tissue sampling	Certain patients at high risk with suspicious nodule morphology, upper lobe location, or both may warrant 12-month follow-up (recommendation 1A).
Multiple				
Low risk [†]	No routine follow-up	CT at 3–6 months, then consider CT at 18–24 months	CT at 3–6 months, then consider CT at 18–24 months	Use most suspicious nodule as guide to management. Follow-up intervals may vary according to size and risk (recommendation 2A).
High risk [†]	Optional CT at 12 months	CT at 3–6 months, then at 18–24 months	CT at 3–6 months, then at 18–24 months	Use most suspicious nodule as guide to management. Follow-up intervals may vary according to size and risk (recommendation 2A).
B: Subsolid Nodules*				
Nodule Type	Size		Comments	
	<6 mm (<100 mm ³)	≥6 mm (>100 mm ³)		
Single				
Ground glass	No routine follow-up	CT at 6–12 months to confirm persistence, then CT every 2 years until 5 years		In certain suspicious nodules < 6 mm, consider follow-up at 2 and 4 years. If solid component(s) or growth develops, consider resection. (Recommendations 3A and 4A).
Part solid	No routine follow-up	CT at 3–6 months to confirm persistence. If unchanged and solid component remains <6 mm, annual CT should be performed for 5 years.		In practice, part-solid nodules cannot be defined as such until ≥6 mm, and nodules <6 mm do not usually require follow-up. Persistent part-solid nodules with solid components ≥6 mm should be considered highly suspicious (recommendations 4A-4C)
Multiple	CT at 3–6 months. If stable, consider CT at 2 and 4 years.	CT at 3–6 months. Subsequent management based on the most suspicious nodule(s).		Multiple <6 mm pure ground-glass nodules are usually benign, but consider follow-up in selected patients at high risk at 2 and 4 years (recommendation 5A).